

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Boo

 Search for

 Limits Preview/Index History Clipboard Details
 Display Show:
☐ 1: [AH001152](#). Bovine preprotach...[gi:163589]

[Links](#)

LOCUS BOVPPTA1 304 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 1''.

ACCESSION M14344

VERSION M14344.1 GI:163580

KEYWORDS .

SEGMENT 1 of 9

SOURCE Bos taurus (cow)

 ORGANISM Bos taurus

 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
 Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 304)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

 MEDLINE [86313713](#)

 PUBMED [3462746](#)

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

 source 1..304
 /organism="Bos taurus"
 /mol_type="genomic DNA"
 /db_xref="taxon:9913"
prim_transcript <162..>304
 /note="preprotachykinin mRNA"
intron 285..>304
 /note="pre-tach-B intron A"

BASE COUNT 63 a 85 c 97 g 59 t

ORIGIN 162 bp upstream of EcoRI site.

 1 ggcagtgggc acagacaggg ccagccgggc tggaggcgcg ggggaccaga cctctcactc
 61 cccggcctgtg atatctccca ggctgtcctt aggagagaaa aatgaatccc ttccctgctg
 121 tttctcccct ccctaagatc tcctcagtgg gctccaaggg ggaattcctg ctcgtcagtg
 181 agcatgtgat tgaactgctg accaaaaatat gccggggtac actggatgcc acgcagatgc
 241 agcttccagt caccgtgact gaggagtaag gccgtgagcc ggggggtgagg gatggcttgg
 301 ggca

//

LOCUS BOVPPTA2 371 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 1'.

ACCESSION M14345

VERSION M14345.1 GI:163581

KEYWORDS .

SEGMENT 2 of 9

SOURCE Bos taurus (cow)

 ORGANISM Bos taurus

 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
 Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 371)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.
 TITLE Structure and gene organization of bovine neuromedin K precursor
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)
 MEDLINE [86313713](#)
 PUBMED [3462746](#)

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.
 FEATURES Location/Qualifiers
 source 1..371
 /organism="Bos taurus"
 /mol_type="genomic DNA"
 /db_xref="taxon:9913"
 prim transcript <1..>371
 /note="preprotachykinin mRNA and introns"
 intron <1..41
 /note="pre-tach-B intron A"
 intron 338..>371
 /note="pre-tach-B intron B"

BASE COUNT 70 a 106 c 107 g 88 t
 ORIGIN 226 bp downstream of exon 1''; 236 bp upstream of Sau3AI site.
 1 gagggcgcta cgtttgctt tctatctgtt ccctgctcca ggttctcagt gaagttcaag
 61 gagggcagct tgacggtcaa ggatcatccag ggccctgggg gtggtgggac tggcaagctg
 121 agcttcaaga agaaggggag tcgttgacct gaggtgactg tacaatgagg aggtgcagcg
 181 gtttcttctt cctggacggg caccggcccc acccccacct gccaccttt gtgggggatc
 241 tcaagtctga acccctgat aggtgtgtgg gtcggagatt gaagaaccct tgaagaggaa
 301 ctgtcttctc caaaccttt ccagaactct ctcaaagtt agcttctctc taccctcggc
 361 ctctttgccc a

//

LOCUS BOVPPTA3 334 bp DNA linear MAM 27-APR-1993
 DEFINITION Bovine preprotachykinin B gene, exon 1.
 ACCESSION M14346
 VERSION M14346.1 GI:163582
 KEYWORDS .
 SEGMENT 3 of 9
 SOURCE Bos taurus (cow)
 ORGANISM Bos taurus
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
 Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 334)
 AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.
 TITLE Structure and gene organization of bovine neuromedin K precursor
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)
 MEDLINE [86313713](#)
 PUBMED [3462746](#)

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.
 FEATURES Location/Qualifiers
 source 1..334
 /organism="Bos taurus"
 /mol_type="genomic DNA"
 /db_xref="taxon:9913"
 intron <1..216
 /note="pre-tach-B intron B"
 prim transcript <17..>334
 /note="preprotachykinin mRNA and introns"
 intron 315..>334
 /note="pre-tach-B intron C"

BASE COUNT 51 a 125 c 84 g 74 t
 ORIGIN About 5700 bp downstream of exon 1'.
 1 tttgtgtgtg gtgtgtgtgt gtgtgtgcgc gtgtgtctga agcctatatc ccccttcacc
 61 ccaccagcc tcccgccccg ccagccaccc tgggattggt gattctcagc cctcccccg

```
121 gctccccag accctcccag agcctttatc agggagctgg gcaggagttc ctgccacatt
181 cccagtgccc tccactccct gtttctctct tcacagcagc agcaccagca gcgttcgtgg
241 ggagcgggca gtgcttcgga ccagctccct gatcctgcta gaccatctgt cccagggtcc
301 aagctgctcc acaggtaggc acaggccaga gaat
```

//

LOCUS BOVPPTA4 179 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 2.

ACCESSION M14347

VERSION M14347.1 GI:163583

KEYWORDS .

SEGMENT 4 of 9

SOURCE Bos taurus (cow)

ORGANISM Bos taurus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 179)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

MEDLINE 86313713PUBMED 3462746

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

source 1..179

/organism="Bos taurus"

/mol_type="genomic DNA"

/db_xref="taxon:9913"

prim transcript <1..>179

/note="preprotachykinin mRNA and introns"

intron <1..40

/note="pre-tach-B intron C"

exon <46..159

/note="preprotachykinin B, (1st expressed exon)"

/number=1

intron 160..>179

/note="pre-tach-B intron D"

BASE COUNT 22 a 58 c 54 g 45 t

ORIGIN 557 bp downstream of exon 1.

1 tcctccctgc ctcccctgtc ctgtctctct gtcctcctag gcatcatgag gagcaccctg

61 ctgtttgcag tcctcctggc cctcagctca gctcggaggt tgggtgcggt ctgtgaggag

121 tcacaggagc aggtggtgcc cgggtgggggt cacagcaagg taaggctccc cctctggtt

//

LOCUS BOVPPTA5 169 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 3.

ACCESSION M14348

VERSION M14348.1 GI:163584

KEYWORDS .

SEGMENT 5 of 9

SOURCE Bos taurus (cow)

ORGANISM Bos taurus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 169)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

MEDLINE 86313713PUBMED 3462746

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

source 1..169
 /organism="Bos taurus"
 /mol_type="genomic DNA"
 /db_xref="taxon:9913"
prim transcript <1..>169
 /note="preprotachykinin mRNA and introns"
intron <1..40
 /note="pre-tach-B intron D"
exon 41..149
 /number=3
intron 150..>169
 /note="pre-tach-B intron E"

BASE COUNT 33 a 54 c 41 g 41 t

ORIGIN About 2000 bp downstream of exon 2.

1 gtgagccttc cttccggccc tctttgtctc tctccccag aaggactcaa atctctacca
 61 gctgccccca tcattgctcc ggagactcta tgatagccgc gtggtctccc tggatggatt
 121 gctcaagatg ctgagcaagg ccagcgtagg taggatatac agcctcagg

//

LOCUS BOVPPTA6 90 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 4.

ACCESSION M14349

VERSION M14349.1 GI:163585

KEYWORDS .

SEGMENT 6 of 9

SOURCE Bos taurus (cow)

ORGANISM Bos taurus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
 Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 90)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

MEDLINE 86313713

PUBMED 3462746

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

source 1..90
 /organism="Bos taurus"
 /mol_type="genomic DNA"
 /db_xref="taxon:9913"
prim transcript <1..>90
 /note="preprotachykinin mRNA and introns"
intron <1..40
 /note="pre-tach-B intron E"
exon 41..70
 /number=4
intron 71..>90
 /note="pre-tach-B intron F"

BASE COUNT 18 a 29 c 16 g 27 t

ORIGIN 195 bp downstream of exon 3.

1 gatcccatc cccagttaca tgcttggtgt tgcttcacag gtcctaagga gtcaccactt
 61 cccagaaac gtgagtagcc tcctttcctt

//

LOCUS BOVPPTA7 114 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine tachykinin B gene, exon 5.

ACCESSION M14350

VERSION M14350.1 GI:163586

KEYWORDS .

SEGMENT 7 of 9

SOURCE Bos taurus (cow)

ORGANISM Bos taurus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 114)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

MEDLINE 86313713

PUBMED 3462746

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

source 1..114
/organism="Bos taurus"
/mol_type="genomic DNA"
/db_xref="taxon:9913" ,

prim transcript <1..>114
/note="preprotachykinin mRNA and introns"

intron <1..40
/note="pre-tach-B intron F"

exon 41..94
/number=5

intron 95..>114
/note="pre-tach-B intron G"

BASE COUNT 22 a 23 c 38 g 31 t

ORIGIN About 480 bp downstream of exon 4.
1 ctcacgtga tgggtgacca tgttgccct ttttatgcag gtgacatgca tgacttcttt
61 gtgggtctca tgggcaagag gaacctccag ccaggtagga gcatgggtggg aggg

//

LOCUS BOVPPTA8 135 bp DNA linear MAM 27-APR-1993

DEFINITION Bovine preprotachykinin B gene, exon 6.

ACCESSION M14351

VERSION M14351.1 GI:163587

KEYWORDS .

SEGMENT 8 of 9

SOURCE Bos taurus (cow)

ORGANISM Bos taurus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.

REFERENCE 1 (bases 1 to 135)

AUTHORS Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.

TITLE Structure and gene organization of bovine neuromedin K precursor

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)

MEDLINE 86313713

PUBMED 3462746

COMMENT Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.

FEATURES Location/Qualifiers

source 1..135
/organism="Bos taurus"
/mol_type="genomic DNA"
/db_xref="taxon:9913"


CDS join(M14347.1:46..159,M14348.1:41..149,M14349.1:41..70,
M14350.1:41..94,41..114)
/note="preprotachykinin B"
/codon_start=1
/protein_id="AAA30723.1"

```

/db_xref="GI:163590"
/translation="MRSTLLFAVILALSSARSLGAVCEESQEQQVPPGGGHSKKDSNLY
QLPPSLLRRLYDSRVVSLDGLLKMLSKASVGPKEsplPQKRDMHDFVGLMGKRNLQP
DTPVDINQENIPSGTfFKYPPSVE"
prim transcript <1..>135
                /note="preprotachykinin mRNA and introns"
intron          <1..40
                /note="pre-tach-B intron G"
exon            41..>114
                /note="preprotachykinin B"
                /number=6
intron          116..>135
                /note="pre-tach-B intron H"
BASE COUNT      41 a      41 c      22 g      31 t
ORIGIN          385 bp downstream of exon 5.
      1 agaaaaatcc ctcctaactt caccctttac atccctccag acactcctgt tgatataaac
     61 caagaaaaca tccccagctt tggcaccttc aagtaccctc caagtgtgga atgaggtaag
    121 gacttactca gggcg
//
LOCUS           BOVPPTA9                      361 bp      DNA      linear      MAM 27-APR-1993
DEFINITION      Bovine preprotachykinin B gene, exon 7.
ACCESSION       M14352
VERSION         M14352.1  GI:163588
KEYWORDS        .
SEGMENT         9 of 9
SOURCE          Bos taurus (cow)
ORGANISM        Bos taurus
                Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
                Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
                Bovidae; Bovinae; Bos.
REFERENCE       1 (bases 1 to 361)
AUTHORS         Kotani,H., Hoshimaru,M., Nawa,H. and Nakanishi,S.
TITLE           Structure and gene organization of bovine neuromedin K precursor
JOURNAL         Proc. Natl. Acad. Sci. U.S.A. 83 (18), 7074-7078 (1986)
MEDLINE         86313713
PUBMED         3462746
COMMENT         Original source text: Bovine intestine, cDNA to mRNA, clone pBNK2.
FEATURES        Location/Qualifiers
                source          1..361
                                /organism="Bos taurus"
                                /mol_type="genomic DNA"
                                /db_xref="taxon:9913"
                prim transcript <1..304
                                /note="preprotachykinin mRNA and introns"
                intron          <1..30
                                /note="pre-tach-B intron H"
BASE COUNT      69 a      103 c      61 g      128 t
ORIGIN          About 2300 bp downstream of exon 6.
      1 tgtagatctg atcttttttt tggctctcag cactccactc ctggactcct gggctacatc
     61 gtgaagacac ccacccttg atgcacatgc ttctatttcc ctctctctc cctaccctg
    121 taacactctt gtgctttgac cccttcttct ctctgtccca ctggctgcag aaactgcctt
    181 gtgagcatcc ccagtcctg taatcattga ctacagatgg ccctggattt tctgtagcat
    241 cctacaaata taatgtctct ctctattcct caacaataaa ggatttttac atattatgac
    301 ctacaggatg tatttggttg gtggatgggt ttttctcttt cttgactctt ttggctgtat
    361 c
//

```

Oct 2 2003 18:31:01



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bio

Search for

Limits

Preview/Index

History

Clipboard

Details

Display

Show:

Send to

Get Subsequence

Feat

☐ 1: NM_009312. Mus musculus tach...[gi:6678208]

Links

LOCUS NM_009312 851 bp mRNA linear ROD 06-OCT-2003

DEFINITION Mus musculus tachykinin 2 (Tac2), mRNA.

ACCESSION NM_009312

VERSION NM_009312.1 GI:6678208

KEYWORDS .

SOURCE Mus musculus (house mouse)

ORGANISM Mus musculus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.

REFERENCE 1 (bases 1 to 851)

AUTHORS Strausberg,R.L., Feingold,E.A., Grouse,L.H., Derge,J.G.,
Klausner,R.D., Collins,F.S., Wagner,L., Shenmen,C.M., Schuler,G.D.,
Altschul,S.F., Zeeberg,B., Buetow,K.H., Schaefer,C.F., Bhat,N.K.,
Hopkins,R.F., Jordan,H., Moore,T., Max,S.I., Wang,J., Hsieh,F.,
Diatchenko,L., Marusina,K., Farmer,A.A., Rubin,G.M., Hong,L.,
Stapleton,M., Soares,M.B., Bonaldo,M.F., Casavant,T.L.,
Scheetz,T.E., Brownstein,M.J., Usdin,T.B., Toshiyuki,S.,
Carninci,P., Prange,C., Raha,S.S., Loquellano,N.A., Peters,G.J.,
Abramson,R.D., Mullahy,S.J., Bosak,S.A., McEwan,P.J.,
McKernan,K.J., Malek,J.A., Gunaratne,P.H., Richards,S.,
Worley,K.C., Hale,S., Garcia,A.M., Gay,L.J., Hulyk,S.W.,
Villalón,D.K., Muzny,D.M., Sodergren,E.J., Lu,X., Gibbs,R.A.,
Fahey,J., Helton,E., Kettelman,M., Madan,A., Rodrigues,S.,
Sanchez,A., Whiting,M., Madan,A., Young,A.C., Shevchenko,Y.,
Bouffard,G.G., Blakesley,R.W., Touchman,J.W., Green,E.D.,
Dickson,M.C., Rodriguez,A.C., Grimwood,J., Schmutz,J., Myers,R.M.,
Butterfield,Y.S., Krzywinski,M.I., Skalska,U., Smailus,D.E.,
Schnerch,A., Schein,J.E., Jones,S.J. and Marra,M.A.

TITLE Generation and initial analysis of more than 15,000 full-length
human and mouse cDNA sequences

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 99 (26), 16899-16903 (2002)

MEDLINE 22388257

PUBMED 12477932

REFERENCE 2 (bases 1 to 851)

AUTHORS Okazaki,Y., Furuno,M., Kasukawa,T., Adachi,J., Bono,H., Kondo,S.,
Nikaido,I., Osato,N., Saito,R., Suzuki,H., Yamanaka,I.,
Kiyosawa,H., Yagi,K., Tomaru,Y., Hasegawa,Y., Nogami,A.,
Schonbach,C., Gojobori,T., Baldarelli,R., Hill,D.P., Bult,C.,
Hume,D.A., Quackenbush,J., Schriml,L.M., Kanapin,A., Matsuda,H.,
Batalov,S., Beisel,K.W., Blake,J.A., Bradt,D., Brusic,V.,
Chothia,C., Corbani,L.E., Cousins,S., Dalla,E., Dragani,T.A.,
Fletcher,C.F., Forrest,A., Frazer,K.S., Gaasterland,T.,
Gariboldi,M., Gissi,C., Godzik,A., Gough,J., Grimmond,S.,
Gustincich,S., Hirokawa,N., Jackson,I.J., Jarvis,E.D., Kanai,A.,
Kawaji,H., Kawasawa,Y., Kedzierski,R.M., King,B.L., Konagaya,A.,
Kurochkin,I.V., Lee,Y., Lenhard,B., Lyons,P.A., Maglott,D.R.,
Maltais,L., Marchionni,L., McKenzie,L., Miki,H., Nagashima,T.,
Numata,K., Okido,T., Pavan,W.J., Pertea,G., Pesole,G.,

*linked Ref
unavailable*

Petrovsky,N., Pillai,R., Pontius,J.U., Qi,D., Ramachandran,S., Ravasi,T., Reed,J.C., Reed,D.J., Reid,J., Ring,B.Z., Ringwald,M., Sandelin,A., Schneider,C., Semple,C.A., Setou,M., Shimada,K., Sultana,R., Takenaka,Y., Taylor,M.S., Teasdale,R.D., Tomita,M., Verardo,R., Wagner,L., Wahlestedt,C., Wang,Y., Watanabe,Y., Wells,C., Wilming,L.G., Wynshaw-Boris,A., Yanagisawa,M., Yang,I., Yang,L., Yuan,Z., Zavolan,M., Zhu,Y., Zimmer,A., Carninci,P., Hayatsu,N., Hirozane-Kishikawa,T., Konno,H., Nakamura,M., Sakazume,N., Sato,K., Shiraki,T., Waki,K., Kawai,J., Aizawa,K., Arakawa,T., Fukuda,S., Hara,A., Hashizume,W., Imotani,K., Ishii,Y., Itoh,M., Kagawa,I., Miyazaki,A., Sakai,K., Sasaki,D., Shibata,K., Shinagawa,A., Yasunishi,A., Yoshino,M., Waterston,R., Lander,E.S., Rogers,J., Birney,E. and Hayashizaki,Y.

TITLE Analysis of the mouse transcriptome based on functional annotation of 60,770 full-length cDNAs

JOURNAL Nature 420 (6915), 563-573 (2002)

MEDLINE [22354683](#)

PUBMED [12466851](#)

REFERENCE 3 (bases 1 to 851)

AUTHORS Kako,K., Munekata,E., Hosaka,M., Murakami,K. and Nakayama,K.

TITLE Cloning and sequence analysis of mouse cDNAs encoding preprotachykinin A and B

JOURNAL Biomed. Res. 14, 253-259 (1993)

COMMENT PROVISIONAL [REFSEQ](#): This record has not yet been subject to final NCBI review. The reference sequence was derived from [D14423.1](#).

FEATURES

source	Location/Qualifiers
	1..851
	/organism="Mus musculus"
	/mol_type="mRNA"
	/strain="ICR"
	/db_xref="taxon:10090"
	/chromosome="10"
gene	1..851
	/gene="Tac2"
	/db_xref="GeneID:21334"
	/db_xref="LocusID:21334"
	/db_xref="MGI:98476"
CDS	265..615
	/gene="Tac2"
	/note="substance K; neurokinin 2; neurokinin A; neuromedin L; neuropeptide K; neurokinin alpha;
	go_component: extracellular space [goid 0005615] [evidence TAS] [pmid 12466851];
	go_function: tachykinin [goid 0008648] [evidence IEA];
	go_function: receptor binding [goid 0005102] [evidence IEA];
	go_process: tachykinin signaling pathway [goid 0007217] [evidence IEA];
	go_process: neuropeptide signaling pathway [goid 0007218] [evidence IEA];
	go_process: synaptic transmission [goid 0007268] [evidence IEA]"
	/codon_start=1
	/product="tachykinin 2"
	/protein_id="NP_033338.1"
	/db_xref="GI:6678209"
	/db_xref="GeneID:21334"
	/db_xref="LocusID:21334"
	/db_xref="MGI:98476"
	/translation="MRSAMLFAAVLALSLAWTFGAVCEEPQEQGRLSKDSLDLYQLPP

```

misc feature      265..429
                  /gene="Tac2"
                  /note="Neurokinin_B; Region: Neurokinin B"
                  /db_xref="CDD:pfam03823"
sig_peptide      265..333
                  /gene="Tac2"
mat_peptide      334..612
                  /gene="Tac2"
                  /product="protachykinin B"
mat_peptide      508..537
                  /gene="Tac2"
                  /product="neurokinin B"
polyA_signal     831..836
                  /gene="Tac2"
polyA_site       851
                  /gene="Tac2"

```

BASE COUNT 199 a 242 c 208 g 202 t

ORIGIN


```

   1  tgactggccc tctctgagtt ccaattcgga tgctggctct ttctccttcc ttcagtctct
  61  ctctctctcc tgggctaaga aacctggaga aagccccttc atctgtgtgg gatgtaaagg
 121  agggctggaa gggtaaaaaa agatgcccgga gctttgaggg gaaaaggaag caatcaagag
 181  acgagggtgc aaagcagagc aagcatcagc cagctccctg atcctgtcag ctatccagct
 241  ccacagcttt gtccttcagg caccatgagg agcgccatgc tgtttgcggc tgcctcggc
 301  ctgagcttgg cttggacctt cggggctgtg tgtgaggagc cacaggagca gggagggagg
 361  ctgagtaagg actctgatct ctatcagctg cctccgtccc tgcttcggag actctacgac
 421  agccgccctg tctctctgga aggattgctg aaagtgctga gcaaggctag cgtgggacca
 481  aaggagacat cacttcacac gaaacgtgac atgcacgact tctttgtggg acttatgggc
 541  aagaggaaca gccaaccaga cactcccacc gacgtggttg aagagaacac ccccagcttt
 601  ggcacacctc aataattccc cagtacagaa aggcactcta cacctggacc ccggattgca
 661  tcataaaaacg tgctgtctcc tgcttcctct tctcctcccc tgctgtgaag gtcctcctgt
 721  tggctccctt ccctactctg catagaagct gcatatgaac agcctcaacc ccatagcaat
 781  tatggtttct gtagtgtcct gcattaaaaa tgctatctct cttcgtcaac aataaagggt
 841  ttttacaag g
```

//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 2 2003 18:31:01



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display Show: Sort Send to

Text Version

☐ 1: Brain Res. 1987 Sep;388(3):243-9.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

A cDNA encoding the precursor of the rat neuropeptide, neurokinin B.

Bonner TI, Affolter HU, Young AC, Young WS 3rd.

Laboratory of Cell Biology, National Institute of Mental Health, Bethesda, M 20892.


We have isolated a cDNA clone from a rat cerebral cortex library which encodes the 116 amino acid precursor of the neuropeptide, neurokinin B. The precursor has 68% amino acid homology to the bovine precursor and encodes a single peptide of the tachykinin family. Except for possible small variations at both ends of the message, there appears to be only a single species of neurokinin B mRNA in rat cerebral cortex. In situ hybridization histochemistry indicates that the message is widely distributed in the rat brain in a pattern distinct from that of substance P message.

PMID: 3479225 [PubMed - indexed for MEDLINE]

Privacy Policy

Display Show: Sort Send to [Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Oct 2 2003 18:01



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

 Search for

Limits

Preview/Index

History

Clipboard

Details

Display

Show:

Send to

☐ 1: [NM_019162](#). *Rattus norvegicus*...[gi:9507174]
[Links](#)


LOCUS NM_019162 727 bp mRNA linear ROD 06-OCT-2003
 DEFINITION *Rattus norvegicus* tachykinin 2 (Tac2), mRNA.
 ACCESSION NM_019162
 VERSION NM_019162.1 GI:9507174
 KEYWORDS .
 SOURCE *Rattus norvegicus* (Norway rat)
 ORGANISM [Rattus norvegicus](#)
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae;
 Rattus.
 REFERENCE 1 (bases 1 to 727)
 AUTHORS Bonner,T.I., Affolter,H.U., Young,A.C. and Young,W.S. III.
 TITLE A cDNA encoding the precursor of the rat neuropeptide neurokinin B
 JOURNAL Brain Res. Mol. Brain Res. 2, 243-249 (1987)
 COMMENT PROVISIONAL REFSEQ: This record has not yet been subject to final
 NCBI review. The reference sequence was derived from [M16410.1](#).
 FEATURES
 source Location/Qualifiers
 1..727
 /organism="Rattus norvegicus"
 /mol_type="mRNA"
 /db_xref="taxon:10116"
 /chromosome="7"
 /map="7q22"
 gene 1..727
 /gene="Tac2"
 /db_xref="GeneID:29191"
 /db_xref="LocusID:29191"
 /db_xref="RGD:3809"
 CDS 115..465
 /gene="Tac2"
 /note="go_function: tachykinin [goid 0008648] [evidence
 IEA];
 go_process: tachykinin signaling pathway [goid 0007217]
 [evidence IEA];
 go_process: synaptic transmission [goid 0007268] [evidence
 IEA]"
 /codon_start=1
 /product="tachykinin 2"
 /protein_id="NP_062035.1"
 /db_xref="GI:9507175"
 /db_xref="GeneID:29191"
 /db_xref="LocusID:29191"
 /db_xref="RGD:3809"
 /translation="MRSAMLFAAVLALSLAWTFGAACEEPQEQGGRLSKSDLSLLPP
 PLLRLYDSRSISLEGLLVLSKASVGPKETSPLQKRDMDHDFVGLMGKRNSQPDTPA
 DVVEENTPSFGVLK"
 misc feature 115..201
 /gene="Tac2"

```

      /note="Neurokinin_B; Region: Neurokinin B"
      /db_xref="CDD:pfam03823"
sig_peptide 115..162
      /gene="Tac2"
mat_peptide 358..387
      /gene="Tac2"
      /product="neurokinin B"
BASE COUNT 155 a 228 c 179 g 165 t
ORIGIN
1  agctgggact ggttcctgcc acctccccga gcccctgcac tcttgtctct ctctgtctcc
61 acagcagcgt caccggctcc cggatcctgc cagccgtcca gctctacagg caccatgagg
121 agcgccatgc tgttcgcggc tgtcctcgcc ctacagcttg catggacctt cggggctgcg
181 tgtgaggaac ctcaggagca gggtagggagg ctacagtaagg actctgatct ctctctgcta
241 cctccacccc tgcttcggag actctacgac agccgctcca tctctctgga aggattgctg
301 aaagtgcgtg gcaaggctag cgtgggaccg aaggagacat cacttccaca gaaacgtgac
361 atgcacgact tctttgtggg acttatgggc aagaggaaca gcccaaccaga cactcccgct
421 gatgtgggtg aagagaacac ccccagcttt ggcgtcctca aataggccag cagtgcagaa
481 aagcactcca ctctcagacc ctggactgca tcataaagac agggttcctg tggcgggtccc
541 agtgccctgc ctccctgctt cctgcctgca aggtcctcct gttggctccc ttcctactc
601 tgcacagatg ctgcatatga acagcctcta ccccatatc aattatggtt tctgtagtgt
661 cctgcattaa aaataccatg tctcctcctc aacaataaag ggtttttaca atggagtgc
721 tgaaaag
//
```

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 2 2003 18:31:01

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by NCSC US Mirror sites: Bolivia Canada China Korea Switzerland Taiwan				
Search <input type="text" value="Swiss-Prot/TrEMBL"/>		<input type="text" value="q9uhf0"/>	<input type="button" value="Go"/>	<input type="button" value="Clear"/>

NiceProt

View of

Swiss-Prot:

Q9UHF0

[Printer-friendly view](#)
[Submit update](#)
[Quick BlastP search](#)

[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)
[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

Note: most headings are clickable, even if they don't appear as links. They link to the [user manual](#) or [other documents](#).

Entry information

Entry name **TKNK_HUMAN**
 Primary accession number **Q9UHF0**
 Secondary accession numbers None
 Entered in Swiss-Prot in Release 40, October 2001
 Sequence was last modified in Release 40, October 2001
 Annotations were last modified in Release 42, September 2003

Name and origin of the protein

Protein name **Neurokinin B [Precursor]**
 Synonyms **NKB**
Neuromedin K
ZNEUROK1
 Gene name **TAC3**
 From **Homo sapiens (Human) [TaxID: 9606]**
 Taxonomy **[Eukaryota](#); [Metazoa](#); [Chordata](#); [Craniata](#); [Vertebrata](#); [Euteleostomi](#); [Mammalia](#); [Eutheria](#); [Primates](#); [Catarrhini](#); [Hominidae](#); [Homo](#).**

References

- [1] SEQUENCE FROM NUCLEIC ACID.
[Sheppard P.](#), [Jelinek L.](#), [Whitmore T.](#), [Blumberg H.](#), [Lehner J.](#), [O'Hara P.](#);
 Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
- [2] SEQUENCE FROM NUCLEIC ACID.
TISSUE=[Placenta](#);
 MEDLINE=20322570; PubMed=10866201; [NCBI, ExPASy, EBI, Israel, Japan]
[Page N.M.](#), [Woods R.J.](#), [Gardiner S.M.](#), [Lomthiasong K.](#), [Gladwell R.T.](#), [Butlin D.J.](#), [Manyonda I.T.](#), [Lowry P.J.](#);
 "Excessive placental neurokinin B secretion during the third trimester causes pre-eclampsia.";
 Nature 405:797-800(2000).
- [3] SEQUENCE FROM NUCLEIC ACID.
TISSUE=[Brain](#);
 MEDLINE=22388257; PubMed=12477932; [NCBI, ExPASy, EBI, Israel, Japan]
[Strausberg R.L.](#), [Feingold E.A.](#), [Grouse L.H.](#), [Derge J.G.](#), [Klausner R.D.](#), [Collins F.S.](#), [Wagner L.](#),

Shenmen C.M., Schuler G.D., Altschul S.F., Zeeberg B., Buetow K.H., Schaefer C.F., Bhat N.K., Hopkins R.F., Jordan H., Moore T., Max S.I., Wang J., Hsieh F., Diatchenko L., Marusina K., Farmer A.A., Rubin G.M., Hong L., Stapleton M., Soares M.B., Bonaldo M.F., Casavant T.L., Scheetz T.E., Brownstein M.J., Usdin T.B., Toshiyuki S., Carninci P., Prange C., Raha S.S., Loquellano N.A., Peters G.J., Abramson R.D., Mullahy S.J., Bosak S.A., McEwan P.J., McKernan K.J., Malek J.A., Gunaratne P.H., Richards S., Worley K.C., Hale S., Garcia A.M., Gay L.J., Hulyk S.W., Villalon D.K., Muzny D.M., Sodergren E.J., Lu X., Gibbs R.A., Fahey J., Helton E., Ketteman M., Madan A., Rodrigues S., Sanchez A., Whiting M., Madan A., Young A.C., Shevchenko Y., Bouffard G.G., Blakesley R.W., Touchman J.W., Green E.D., Dickson M.C., Rodriguez A.C., Grimwood J., Schmutz J., Myers R.M., Butterfield Y.S.N., Krzywinski M.I., Skalska U., Smailus D.E., Schnerch A., Schein J.E., Jones S.J.M., Marra M.A.;

"Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences.";

Proc. Natl. Acad. Sci. U.S.A. 99:16899-16903(2002).

Comments

- **FUNCTION:** Tachykinins are active peptides which excite neurons, evoke behavioral responses, are potent vasodilators and secretagogues, and contract (directly or indirectly) many smooth muscles (*By similarity*).
- **SUBCELLULAR LOCATION:** Secreted.
- **DEVELOPMENTAL STAGE:** In pregnancy, the expression of NKB is confined to the outer syncytiotrophoblast of the placenta, significant concentrations of NKB can be detected in plasma as early as week 9, and plasma concentrations of NKB are grossly elevated in pregnancy-induced hypertension and pre-eclampsia.
- **SIMILARITY:** Belongs to the tachykinin family.

Copyright

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch)


Cross-references

EMBL	AF186112; AAF01430.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence] AF216586; AAF76980.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence] BC032145; AAH32145.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence]
Genew	HGNC:11521 ; TAC3.
CleanEx	HGNC:11521 ; TAC3.
GeneCards	TAC3 .
GeneLynx	TAC3 ; Homo sapiens.
GenAtlas	TAC3 .
MIM	162330 [NCBI / EBI]. GO:0005615 ; Cellular component: extracellular space (<i>traceable author statement</i>). GO:0005625 ; Cellular component: soluble fraction (<i>traceable author statement</i>). GO:0008648 ; Molecular function: tachykinin (<i>traceable author statement</i>). GO:0007565 ; Biological process: pregnancy (<i>traceable author statement</i>). GO:0007217 ; Biological process: tachykinin signaling pathway (<i>traceable author statement</i>).
SOURCE	TAC3 ; Homo sapiens.
Ensembl	Q9UHF0; Homo sapiens. [Entry / Contig view] IPR003635 ; Neurokinin.

InterPro [IPR002040](#); Tachy_Neurokinin.
[Graphical view of domain structure.](#)
 Pfam [PF03823](#); Neurokinin_B; 1.
 ProDom [PD020370](#); Neurokinin; 1.
[\[Domain structure / List of seq. sharing at least 1 domain\]](#)
 PROSITE [PS00267](#); TACHYKININ; 1.
 HOVERGEN [\[Family / Alignment / Tree\]](#)
 BLOCKS [Q9UHF0](#).
 ProtoNet [Q9UHF0](#).
 ProtoMap [Q9UHF0](#).
 PRESAGE [Q9UHF0](#).
 DIP [Q9UHF0](#).
 ModBase [Q9UHF0](#).
 SWISS-2DPAGE [Get region on 2D PAGE.](#)

Keywords

Tachykinin; Neuropeptide; Cleavage on pair of basic residues; Amidation; Signal.

Features

[Feature table viewer](#)



[Feature aligner](#)

Key	From	To	Length	Description
SIGNAL	1	16	16	POTENTIAL.
PROPEP	17	78	62	BY SIMILARITY.
PEPTIDE	81	90	10	NEUROKININ B.
PROPEP	94	121	28	BY SIMILARITY.
MOD_RES	90	90		AMIDATION (G-91 PROVIDE AMIDE GROUP) (BY SIMILARITY).

Sequence information

Length: **121 AA** [This is the length of the unprocessed precursor]
 Molecular weight: **13438 Da** [This is the MW of the unprocessed precursor]
 CRC64: **14C9AFE2EE9EDECA** [This is a checksum on the sequence]

10	20	30	40	50	60
MRIMLLFTAI	LAFSLAQSG	AVCKEPQEEV	VPGGGRSKRD	PDLYQLLQRL	FKSHSSLEGL
70	80	90	100	110	120
LKALSQASTD	PKESTSPEKR	DMHDFVGLM	GKRSVQPDSP	TDVNQENVPS	FGILKYPPRA

E

[Q9UHF0 in FASTA format](#)

[View entry in original Swiss-Prot format](#)

[View entry in raw text format \(no links\)](#)

[Report form for errors/updates in this Swiss-Prot entry](#)

BLAST [BLAST submission on ExPASy/SIB](#)
or at [NCBI \(USA\)](#)




Sequence analysis tools: [ProtParam](#), [ProtScale](#),
[Compute pI/Mw](#), [PeptideMass](#), [PeptideCutter](#),
[Dotlet \(Java\)](#)




[ScanProsite](#), [MotifScan](#)



Search the [SWISS-MODEL Repository](#)

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by NCSC US				
Mirror sites:				
Bolivia	Canada	China	Korea	Switzerland
Taiwan				



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: Gynecol Obstet Invest. 1988;25(4):258-61.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Effects of human calcitonin gene-related peptide and substance on human intracervical arteries.

Hansen V, Schifter S, Allen J, Maigaard S, Forman A.

Department of Obstetrics and Gynecology, University of Aarhus, Denmark.

The contractile effects of substance P and human calcitonin gene-related peptide (human CGRP) on isolated human intracervical arteries were studied. Human cervical tissue specimens were excised after hysterectomy at various phases of the menstrual cycle (n = 14) and small intracervical arteries were dissected free by microtechnique. Ring preparations of the vessels were prepared and mounted in organ baths, and isometric circular tension was recorded. Neither compound affected resting tension. Both peptides showed potent relaxing effects on vessels precontracted by noradrenaline 10(-5) M. Substance P exhibited the higher potency, while human CGRP showed the higher efficacy. The relaxing effects of the two compounds were unaffected by pretreatment with indomethacin 10(-6) M, propranolol 10(-6) M and atropine 10(-6) M. The results support a role for the two peptides in the regulation of cervical blood flow.

PMID: 2456971 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Oct 2 2003 13:4